

Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order INTRAVENOUS IMMUNE GLOBULIN (IVIG)

Patient Name:
Date of Birth:
Med. Rec. No (TVC MRN Only):

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

Anticipated Start Date: Patient to follow up with provider on date: ***This plan will expire after 365 days, unless otherwise specified below*** Orders expire:						
•		Height:	cm			
Allergies:						
_				e:		
GUIDELINES FO	R PRESC	RIBING:				
 In patients considere specified If the prefinterchang Pharmacis Ideal Body IB' IB' If It 	erred brange the proc st will rour y Weight (W Males (W Female neight <60	be at risk of renal ill be infused per L d of IVIG (Priviger duct with another l d dose to the nea IBW) will be used kg) = 50 + (2.3 x (l s (kg) = 45.5 + (2. inches, use 50 kg	Il failure, a decrease in Legacy Institutional Rat n) is unavailable, or on brand. Please specify it to dose IVIG. (height in inches – 60)) a x (height in inches – g (male) and 45.5 kg (fe			
LABS TO BE DR	AWN (or	lers must be plac	ced in TVC Epic by or	dering provider if TVC provider):		
☐ CBC with di	fferential, n, Routine,	Routine, ONCE, e	every(visit)(days (visit)(days)(weeks	(weeks)(months) Circle one s)(weeks)(months) Circle one)(months) Circle one		
PRE-MEDICATION	ONS: (Adn	ninister 30 minutes	s prior to infusion)			
-		select which med appropriate box(s		rou would like the patient to receive prior to		
Acetaminopher 325 mg 500 mg	□ 650	DL) tablet oral, ON) mg)0 mg	ICE, every visit			
Diphenhydrami ☐ 25 mg ☐ 50 mg	ne (BENA	DRYL) tablet oral,	, ONCE, every visit			
Cetirizine (ZYR	TEC) table	et oral, ONCE, eve	ery visit (Choose as al	ternative to diphenhydramine if needed		

Page 1 of 3 Last updated 6/2024



Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order INTRAVENOUS IMMUNE GLOBULIN (IVIG)

Patient Name:	
Date of Birth:	
Med. Rec. No (TVC MRN Only):	

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

MEDICATIONS: (must check at least one):

Intravenous Immune Globulin (IVIG) Privigen 10% (preferred brand): (Pharmacist will round dose to nearest 5 gm vial and modify brand/selection based upon availability during order verification)						
 □ 0.2 g/kg IV, ONCE, every visit □ 0.4 g g/kg IV, ONCE, every visit □ 0.5 g/kg IV, ONCE, every visit □ 1 g/kg IV, ONCE, every visit □ g IV, ONCE (for doses titrated to IgG level), every visit 						
Interval: □ Once □ Daily x doses □ Every weeks for doses □ Other						
Specifications: ☐ Patient requires a specific brand of IVIG (other than listed above) Please specify here: ☐ Patient requires IVIG at a 5% concentration (note: currently not a standard stocked item) Infuse per Legacy Immune Globulin Infusion Rate Guidelines (decrease rate of infusion in patients who may be at risk of renal failure) or specify rate below:						

AS NEEDED MEDICATIONS:

Acetaminophen 650 mg oral, EVERY 4 HOURS AS NEEDED for fever, headache or pain

NURSING ORDERS (TREATMENT PARAMETERS):

- 1. Vital signs, every visit: Assess vital signs before initiating IVIG infusion. During the first two infusions: assess vital signs at 15 minutes, 30 minutes, 1 hour, then hourly for remainder of infusion. For subsequent infusions: if the patient has been stable without adverse reactions, the frequency of vital signs is discretionary
- 2. Nursing communication order, every visit: Manage line per LH policy 904.4007 IV Catheter Insertion (Peripheral) and LH 904.4004 IV Access: Central Catheters
- 3. Nursing communication orders, every visit: Manage hypersensitivity reactions per LH 906.6606

Page 2 of 3 Last updated 6/2024



Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order INTRAVENOUS IMMUNE GLOBULIN (IVIG)

Patient Name:
Date of Birth:
Med. Rec. No (TVC MRN Only):

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

HYPERSENSITIVITY MEDICATIONS: Refer to LH policy 906.6606 Initiation of Emergency Measures for Adult Oncology, Radiation Oncology and Infusion Clinic Patients

- 1. diphenhydrAMINE 25-50 mg IV, AS NEEDED x1 for hypersensitivity reaction
- 2. Famotidine 20 mg IV, AS NEEDED x1 dose for hypersensitivity reaction
- 3. Hydrocortisone 100 mg IV, AS NEEDED x1 dose for hypersensitivity reaction
- 4. EPINEPHrine 0.3 mg IM, AS NEEDED x1 dose for hypersensitivity reaction
- 5. Naloxone (Narcan) 0.4 mg IV, AS NEEDED x1 dose for diminished respiratory rate
- 6. Sodium chloride 0.9% 1000 mL IV, AS NEEDED x 1 dose for alteration in hemodynamic status
- 7. Meperidine 12.5-25 mg IV, AS NEEDED x 2 for infusion-related rigors
- 8. Nursing communication order, every visit: Please follow treatment algorithm for acute infusion reaction

Please check the appropriate box for the patient's prefer	red clinic location:		
□ Legacy Day Treatment Unit – The Vancouver Clinic Building A department of Salmon Creek Medical Center 700 NE 87 th Avenue, Suite 360 Vancouver, WA 98664 Phone number: 360-896-7070 Fax number: 360-487-5773	☐ Legacy Emanuel Day Treatment Unit A department of Emanuel Medical Center 501 N Graham Street, Suite 540 Portland, OR 97227 Phone number: 503-413-4608 Fax number: 503-413-4887		
Legacy Salmon Creek Day Treatment Unit Legacy Salmon Creek Medical Center 2121 NE 139 th Street, Suite 110 Vancouver, WA 98686 Phone number: 360-487-1750 Fax number: 360-487-5773	□ Legacy STEPS Clinic A department of Silverton Medical Center Legacy Woodburn Health Center 1475 Mt Hood Ave Woodburn, OR 97071 Phone number: 503-982-1280 Fax number: 503-225-8723		
Provider signature:	Date/Time:		
Printed Name:	Phone:	Fax:	
Organization/Department:			

Page 3 of 3 Last updated 6/2024