

### Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order RITUXIMAB (RITUXAN) FOR NON-ONCOLOGY INDICATIONS

Patient Name:	
Date of Birth:	
Med. Rec. No (TVC MRN Only):	

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

	expire aft	er 365 days, u	Patient to follow up with provider on date: unless otherwise specified below***
Weight:			
Allergies:		_	
Diagnosis:			
GUIDELINES FO	R PRESC	RIBING:	
2. Hepatitis	B (Hep B s	•	CARD and most recent provider chart or progress note. and core antibody) screening must be completed prior to initiation or be infected
PRE-SCREENIN	G: (Resul	ts must be ava	ailable prior to initiation of therapy)
			ult Date: Desitive / Negative esult Date: Desitive / Negative
LABS TO BE DE	RAWN (or	lers must be p	placed in TVC Epic by ordering provider if TVC provider):
☐ CBC with	differentia	I, Routine, ever	every(visit)(days)(weeks)(months) - Circle one ry(visit)(days)(weeks)(months) - Circle one
☐ acetamin ☐ 65 ☐ 32 ☐ 0 ☐ diphenhy ☐ 25 ☐ 50 ☐ cetirizine	ophen (TY 50 mg 25 mg ther: drAMINE ( mg mg (ZYRTEC)	LENOL) tablet, BENADRYL) ta	utes prior to infusion) oral, every visit  ablet, oral, every visit  very visit (Choose as alternative to diphenhydramine, if needed)
□ 12	EDNISolo 25 mg 2.5 mg	ne sodium succ	cinate (SOLU-MEDROL) IV, every visit

Page 1 of 3 Last updated 6/2024



# Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order RITUXIMAB (RITUXAN) FOR NON-ONCOLOGY INDICATIONS

Patient Name:
Date of Birth:
Med. Rec. No (TVC MRN Only):

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

	CATIONS: (must check one): se: (Pharmacist will use most recent weight at initiation and round dose to the nearest whole vial)
	riTUXimab (RITUXAN) 375 mg/m2 = mg IV in NaCl 0.9% to a final concentration of 2 mg/mL. First infusion or prior infusion reactions: start at 50 mg/hr x 30 min, then may increase by 50 mg/hr every 30 minutes if tolerated (NTE 400 mg/hr). Subsequent infusions if no reactions infuse rituximab at 100 mg/hr for the first 30 min. If no infusion related reactions are seen, increase rate by 100 mg/hour every 30 minutes to a maximum of 400 mg/hr
	<b>riTUXimab (RITUXAN) mg IV in NaCl 0.9%</b> to a final concentration 2 mg/mL. First infusion or prior infusion reactions: start at 50 mg/hr x 30 min, then may increase by 50 mg/hr every 30 minutes if tolerated (NTE 400 mg/hr). Subsequent infusions if no reactions infuse rituximab at 100 mg/hr for the first 30 min. If no infusion related reactions are seen, increase rate by 100 mg/hour every 30 minutes to a maximum of 400 mg/hr
Interv	al:
	Once Every 2 weeks x 2 doses Every weeks x doses Weekly x 4 doses Other

#### **AS NEEDED MEDICATIONS:**

- acetaminophen 650 mg oral, EVERY 4 HOURS AS NEEDED for fever, headache or pain
- diphenhydrAMINE 25 mg oral, EVERY 4 HOURS AS NEEDED for itching
- meperidine 25-50 mg IV, EVERY 2 HOURS AS NEEDED (NTE 50 mg/hr) for rigors in the absence of hypotension

### **NURSING ORDERS (TREATMENT PARAMETERS):**

- 1. Vital signs, every visit: Monitor and record vital signs, tolerance, and presence of infusion-related reactions prior to infusion and at the end of infusion.
- 2. Nursing communication order, every visit: Manage line per LH policy 904.4007 IV Catheter Insertion (Peripheral) and LH 904.4004 IV Access: Central Catheters.
- 3. Nursing communication orders, every visit: Manage hypersensitivity reactions per LH 906.6606.

Page 2 of 3 Last updated 6/2024



## Legacy Day Treatment Unit Provider's Orders

Adult Ambulatory Infusion Order RITUXIMAB (RITUXAN) FOR NON-ONCOLOGY INDICATIONS

Patient Name:
Date of Birth:
Med. Rec. No (TVC MRN Only):

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

**HYPERSENSITIVITY MEDICATIONS:** Refer to LH policy 906.6606 Initiation of Emergency Measures for Adult Oncology, Radiation Oncology and Infusion Clinic Patients

- 1. hydrocortisone 100 mg IV, AS NEEDED x1 dose for hypersensitivity reaction
- 2. diphenhydrAMINE 25-50 mg IV, EVERY 2 HOURS AS NEEDED for hypersensitivity reaction (Max dose: 50 mg)
- 3. famotidine 20 mg IV, AS NEEDED x1 dose for hypersensitivity reaction
- 4. EPINEPHrine 0.3 mg IM, AS NEEDED x1 dose for hypersensitivity reaction
- 5. naloxone (Narcan) 0.4 mg IV, AS NEEDED x1 dose for diminished respiratory rate
- 6. sodium chloride 0.9% 1000 mL IV, AS NEEDED x 1 dose for alteration in hemodynamic status
- 7. Nursing communication order, every visit: Please follow treatment algorithm for acute infusion reaction.

Please check the appropriate box for the patient's preferred clinic location: ☐ Legacy Day Treatment Unit – ☐ Legacy Emanuel Day Treatment Unit The Vancouver Clinic Building A department of Emanuel Medical Center A department of Salmon Creek Medical Center 501 N Graham Street, Suite 540 700 NE 87<sup>th</sup> Avenue, Suite 360 Portland, OR 97227 Vancouver, WA 98664 Phone number: 503-413-4608 Phone number: 360-896-7070 Fax number: 503-413-4887 Fax number: 360-487-5773 ☐ Legacy Salmon Creek Day Treatment Unit ☐ Legacy STEPS Clinic Legacy Salmon Creek Medical Center A department of Silverton Medical Center 2121 NE 139th Street, Suite 110 Legacy Woodburn Health Center Vancouver, WA 98686 1475 Mt Hood Ave Phone number: 360-487-1750 Woodburn, OR 97071 Fax number: 360-487-5773 Phone number: 503-982-1280 Fax number: 503-225-8723 Provider signature: \_\_\_\_\_ Date/Time: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_ Organization/Department:

Page 3 of 3 Last updated 6/2024