 <p>LEGACY HEALTH</p>	<p>Legacy Day Treatment Unit Provider's Orders</p> <p>Adult Ambulatory Infusion Order VEDOLIZUMAB (ENTYVIO) INFUSION</p>	<p>Patient Name: _____</p> <p>Date of Birth: _____</p> <p>Med. Rec. No (TVC MRN Only): _____</p>
<p>ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE</p>		

Anticipated Start Date: _____ **Patient to follow up with provider on date:** _____

*****This plan will expire after 365 days, unless otherwise specified below*****

Orders expire: _____

Weight: _____ kg **Height:** _____ cm

Allergies: _____

Diagnosis: _____

Diagnosis Code: _____ (please include primary and secondary diagnosis codes)

GUIDELINES FOR PRESCRIBING:

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Hepatitis B (Hep B surface antigen and core antibody total) screening must be completed prior to initiation of treatment and the patient should not be infected. Please send results with order.
3. A Tuberculin test must have been placed and read as negative prior to initiation of treatment (PPD or QuantiFERON Gold blood test). Please send results with order. If result is indeterminate, a follow up chest X-ray must be performed to rule out TB. Please send results with order.
4. Patients should not have an active ongoing infection, signs or symptoms of malignancy, or moderate to severe heart failure at the onset of therapy. Baseline liver function tests should be normal. 2. Patient should have regular monitoring for infection, malignancy, and liver abnormalities throughout therapy.

PRE-SCREENING (orders must be placed in TVC Epic by ordering provider if TVC provider):


- Hepatitis B surface antigen and core antibody total test results scanned with orders.
- Tuberculin skin test or QuantiFERON Gold blood test results scanned with orders.
- Chest X-Ray result scanned with orders if TB test result is indeterminate.

LABS (orders must be placed in TVC Epic by ordering provider if TVC provider):

- Complete Metabolic Panel, Routine, ONCE, every visit
- CBC with differential, Routine, ONCE, every visit

NURSING ORDERS (TREATMENT PARAMETERS):

1. TREATMENT PARAMETER – Hold treatment and contact provider if Hepatitis B surface antigen or core antibody total test result is positive, TB test result is positive, or if screening has not been performed.
2. TREATMENT PARAMETER – Hold treatment and contact provider if patient has signs or symptoms of infection.
3. VITAL SIGNS – Monitor patient for signs and symptoms of hypersensitivity during the infusion.
4. Nursing communication order, every visit: Manage line per LH policy 904.4007 IV Catheter Insertion (Peripheral) and LH 904.4004 IV Access: Central Catheters.
5. Nursing communication order, every visit: Manage hypersensitivity reactions per LH policy 906.6606.

 <p>LEGACY HEALTH</p>	<p>Legacy Day Treatment Unit Provider's Orders</p> <p>Adult Ambulatory Infusion Order VEDOLIZUMAB (ENTYVIO) INFUSION</p>	<p>Patient Name: _____</p> <p>Date of Birth: _____</p> <p>Med. Rec. No (TVC MRN Only): _____</p>
<p>ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE</p>		

MEDICATIONS:

- vedolizumab (ENTYVIO) 300 mg in sodium chloride 0.9%, intravenous, ONCE over 30 minutes
Interval (must check at least one)
 - Initial dosing:** on week 0, 2, and 6
 - Maintenance dosing:** every 8 weeks thereafter
 - Other:** _____

AS NEEDED MEDICATIONS

- acetaminophen (TYLENOL) tablet, 650 mg, oral, ONCE, every visit
- diphenhydrAMINE (BENADRYL) capsule, 50 mg, oral, ONCE, every visit.

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION - If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Refer to LH policy 906.6606 Initiation of Emergency Measures for Adult Oncology, Radiation Oncology and Infusion Clinic Patients
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

Please check the appropriate box for the patient's preferred clinic location:

- | | |
|---|---|
| <input type="checkbox"/> Legacy Day Treatment Unit –
The Vancouver Clinic Building
<i>A department of Salmon Creek Medical Center</i>
700 NE 87 th Avenue, Suite 360
Vancouver, WA 98664
Phone number: 360-896-7070
Fax number: 360-487-5773 | <input type="checkbox"/> Legacy Emanuel Day Treatment Unit
<i>A department of Emanuel Medical Center</i>
501 N Graham Street, Suite 540
Portland, OR 97227
Phone number: 503-413-4608
Fax number: 503-413-4887 |
| <input type="checkbox"/> Legacy Salmon Creek Day Treatment Unit
Legacy Salmon Creek Medical Center
2121 NE 139 th Street, Suite 110
Vancouver, WA 98686
Phone number: 360-487-1750
Fax number: 360-487-5773 | <input type="checkbox"/> Legacy STEPS Clinic
<i>A department of Silverton Medical Center</i>
Legacy Woodburn Health Center
1475 Mt Hood Ave
Woodburn, OR 97071
Phone number: 503-982-1280
Fax number: 503-225-8723 |

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Organization/Department: _____